



Available at  
**www.ElsevierMathematics.com**  
POWERED BY SCIENCE @ DIRECT®

J. Math. Anal. Appl. 288 (2003) 708–716

*Journal of*  
MATHEMATICAL  
ANALYSIS AND  
APPLICATIONS

www.elsevier.com/locate/jmaa

## Three positive solutions of a nonlinear three-point boundary value problem

Jian-Ping Sun,<sup>a,\*</sup> Wan-Tong Li,<sup>b,2</sup> and Ya-Hong Zhao<sup>a</sup>

<sup>a</sup> Department of Applied Mathematics, Lanzhou University of Technology, Lanzhou, Gansu 730050, People's Republic of China

<sup>b</sup> Department of Mathematics, Lanzhou University, Lanzhou, Gansu 730000, People's Republic of China

Received 9 May 2003

Submitted by J. Henderson

---

### Abstract

In this paper, existence criteria for three positive solutions of the nonlinear three-point boundary value problem

$$\begin{cases} u''(t) + a(t)u'(t) + b(t)u(t) + h(t)f(u) = 0, & t \in (0, 1), \\ u(0) = 0, & u(1) = \alpha u(\eta), \end{cases}$$

are established by using the Leggett–Williams fixed point theorem. An example is also included to illustrate the importance of the result obtained.

© 2003 Elsevier Inc. All rights reserved.

**Keywords:** Multi-point boundary value problem; Positive solution; Fixed point

---

---

\* Corresponding author.

E-mail addresses: sunjianping9386@sina.com (J.-P. Sun), wtli@lzu.edu.cn (W.-T. Li).

<sup>1</sup> Partially supported by the Key Research and Development Program for Outstanding Groups of Lanzhou University of Technology.

<sup>2</sup> Partially supported by the NNSF of China (10171040), the NSF of Gansu Province of China (ZS011-A25-007-Z), the Foundation for University Key Teacher by the Ministry of Education of China, and the Teaching and Research Award Program for Outstanding Young Teachers in Higher Education Institutions of Ministry of Education of China.

## 1. Introduction

Recently, multi-point boundary value problems have received many attentions from many authors [4–6,8–11]. In particular, Ma and Wang [10] obtained the existence of one positive solution for more general three-point boundary value problem

$$u''(t) + a(t)u'(t) + b(t)u(t) + h(t)f(u) = 0, \quad t \in (0, 1), \quad (1.1)$$

$$u(0) = 0, \quad u(1) = \alpha u(\eta), \quad 0 < \eta < 1, \quad (1.2)$$

under the assumption that  $f$  is either superlinear or sublinear, where (H1)  $f \in C([0, \infty), [0, \infty))$ ; (H2)  $h \in C([0, 1], [0, \infty))$  and there exists  $x_0 \in [0, 1]$  such that  $h(x_0) > 0$ ; (H3)  $a \in C[0, 1]$ ,  $b \in C([0, 1], (-\infty, 0))$ ; (H4)  $0 < \alpha\phi_1(\eta) < 1$ , where  $\phi_1$  is the unique solution of the linear boundary value problem

$$\begin{cases} \phi_1''(t) + a(t)\phi_1'(t) + b(t)\phi_1(t) = 0, & t \in (0, 1), \\ \phi_1(0) = 0, & \phi_1(1) = 1. \end{cases} \quad (1.3)$$

Their main tool is a fixed point theorem for mapping defined on Banach spaces with cones [1,3]. For background information and applications of such a fixed point theorem, one can refer to [2,8,12].

In this paper we will consider the existence of at least three positive solutions for the above boundary value problems. Our tool is the well-known Leggett–Williams fixed point theorem [7].

In order to obtain our main results, we need the following concepts and Leggett–Williams fixed point theorem.

Let  $E$  be a real Banach space with cone  $P$ . A map  $\alpha: P \rightarrow [0, +\infty)$  is said to be a nonnegative continuous concave functional on  $P$  if  $\alpha$  is continuous and

$$\alpha(tx + (1-t)y) \geq t\alpha(x) + (1-t)\alpha(y)$$

for all  $x, y \in P$  and  $t \in [0, 1]$ . Let  $a_0, b_0$  be two numbers such that  $0 < a_0 < b_0$  and  $\alpha$  a nonnegative continuous concave functional on  $P$ . We define the following convex sets

$$P_{a_0} = \{x \in P: \|x\| < a_0\},$$

$$P(\alpha, a_0, b_0) = \{x \in P: a_0 \leq \alpha(x), \|x\| \leq b_0\}.$$

**Theorem** (Leggett–Williams fixed point theorem). *Let  $A: \bar{P}_c \rightarrow \bar{P}_c$  be completely continuous and  $\alpha$  be a nonnegative continuous concave functional on  $P$  such that  $\alpha(x) \leq \|x\|$  for all  $x \in \bar{P}_c$ . Suppose there exist  $0 < d_0 < a_0 < b_0 \leq c$  such that*

- (i)  $\{x \in P(\alpha, a_0, b_0): \alpha(x) > a_0\} \neq \emptyset$  and  $\alpha(Ax) > a_0$  for  $x \in P(\alpha, a_0, b_0)$ ;
- (ii)  $\|Ax\| < d_0$  for  $\|x\| \leq d_0$ ;
- (iii)  $\alpha(Ax) > a_0$  for  $x \in P(\alpha, a_0, c)$  with  $\|Ax\| > b_0$ .

*Then  $A$  has at least three fixed points  $x_1, x_2, x_3$  in  $\bar{P}_c$  satisfying*

$$\|x_1\| < d_0, \quad a_0 < \alpha(x_2), \quad \|x_3\| > d_0 \quad \text{and} \quad \alpha(x_3) < a_0.$$

## 2. Main results

To state the main result of this paper, we need the following lemma, which was established by Ma and Wang [10].

**Lemma 2.1.** Assume that (H3) holds. Let  $\phi_1$  and  $\phi_2$  be the solutions of Eq. (1.3) and

$$\begin{cases} \phi_2''(t) + a(t)\phi_2'(t) + b(t)\phi_2(t) = 0, & t \in (0, 1), \\ \phi_2(0) = 1, & \phi_2(1) = 0. \end{cases} \quad (2.1)$$

Then

- (i)  $\phi_1$  is strictly increasing on  $[0, 1]$ ;
- (ii)  $\phi_2$  is strictly decreasing on  $[0, 1]$ .

In view of (H2),  $h \in C([0, 1], [0, \infty))$  and there exists  $x_0 \in [0, 1]$  such that  $h(x_0) > 0$ , and hence we may assume that  $x_0 \in (0, 1)$ . Take  $\delta \in (0, 1/2)$  such that  $x_0 \in (\delta, 1 - \delta)$ .

For convenience, we let

$$G(t, s) = \frac{1}{\phi_1'(0)} \begin{cases} \phi_1(t)\phi_2(s), & s \geq t, \\ \phi_1(s)\phi_2(t), & s \leq t, \end{cases}$$

$$D = \max_{t \in [0, 1]} \int_0^1 G(t, s)p(s)h(s)ds + \frac{\alpha}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s)p(s)h(s)ds,$$

and

$$C = \min_{t \in [\delta, 1 - \delta]} \int_{\delta}^{1 - \delta} G(t, s)p(s)h(s)ds + \frac{\alpha\phi_1(\delta)}{1 - \alpha\phi_1(\eta)} \int_{\delta}^{1 - \delta} G(\eta, s)p(s)h(s)ds,$$

where  $p(t) = \exp(\int_0^t a(s)ds)$ .

For the function  $G(t, s)$ , it follows from Lemma 2.1 that

$$0 \leq G(t, s) \leq G(s, s), \quad (t, s) \in [0, 1] \times [0, 1], \quad (2.2)$$

and

$$G(t, s) \geq \gamma G(s, s), \quad (t, s) \in [\delta, 1 - \delta] \times [0, 1], \quad (2.3)$$

where  $\gamma = \min\{\phi_1(\delta), \phi_2(1 - \delta)\}$ .

Our main result is the following theorem.

**Theorem 2.1.** Assume that (H1)–(H4) hold and that there exist numbers  $d_1$  and  $d_0$  with  $0 < d_0 < d_1$  such that

$$f(u) < \frac{d_0}{D}, \quad u \in [0, d_0], \quad (2.4)$$

and

$$f(u) > d_1/C, \quad u \in \left[d_1, \frac{d_1}{\gamma}\right]. \quad (2.5)$$

Then the boundary value problem (1.1)–(1.2) has at least three positive solutions if the following condition holds:

(H5) There exists a number  $c > d_1/\gamma$  such that  $f(u) < c/D$  for  $u \in [0, c]$ .

**Proof.** Let  $E$  be the set  $C[0, 1]$  of all real continuous functions defined on  $[0, 1]$  endowed with the usual linear structure and the maximum norm. Set

$$P = \{u \in E: u(t) \geq 0, t \in [0, 1]\}.$$

Then it is easily seen that  $P$  is a cone in  $E$ . For  $u \in P$ , define

$$\alpha(u) = \min_{t \in [\delta, 1-\delta]} u(t)$$

and

$$\begin{aligned} (Au)(t) &= \int_0^1 G(t, s)p(s)h(s)f(u(s))ds \\ &\quad + \frac{\alpha\phi_1(t)}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s)p(s)h(s)f(u(s))ds, \quad t \in [0, 1]. \end{aligned}$$

It is easy to check that  $\alpha$  is a nonnegative continuous concave functional on  $P$  with  $\alpha(x) \leq \|x\|$  for  $x \in P$  and that [10]  $A: P \rightarrow P$  is completely continuous and fixed points of  $A$  are solutions of the boundary value problem (1.1)–(1.2).

We first assert that if there exists a positive number  $r$  such that  $f(u) < r/D$  for  $u \in [0, r]$ , then  $A: \bar{P}_r \rightarrow P_r$ .

Indeed, if  $u \in \bar{P}_r$ , then, in view of Lemma 2.1, for  $t \in [0, 1]$ ,

$$\begin{aligned} (Au)(t) &= \int_0^1 G(t, s)p(s)h(s)f(u(s))ds \\ &\quad + \frac{\alpha\phi_1(t)}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s)p(s)h(s)f(u(s))ds \\ &< \frac{r}{D} \left[ \int_0^1 G(t, s)p(s)h(s)ds + \frac{\alpha\phi_1(t)}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s)p(s)h(s)ds \right] \\ &\leq \frac{r}{D} \left[ \max_{t \in [0, 1]} \int_0^1 G(t, s)p(s)h(s)ds + \frac{\alpha}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s)p(s)h(s)ds \right] \\ &= r. \end{aligned}$$

Thus,  $\|Au\| < r$ , i.e.,  $Au \in P_r$ .

Hence, we have shown that if (2.4) and (H5) hold, then  $A$  maps  $\bar{P}_{d_0}$  into  $P_{d_0}$  and  $\bar{P}_c$  into  $P_c$ .

Next, we assert that  $\{u \in P(\alpha, d_1, d_1/\gamma): \alpha(u) > d_1\} \neq \emptyset$  and  $\alpha(Au) > d_1$  for all  $u \in P(\alpha, d_1, d_1/\gamma)$ .

In fact, the constant function

$$\frac{d_1 + d_1/\gamma}{2} \in \{u \in P(\alpha, d_1, d_1/\gamma): \alpha(u) > d_1\}.$$

Moreover, for  $u \in P(\alpha, d_1, d_1/\gamma)$ , we have

$$d_1/\gamma \geq \|u\| \geq u(t) \geq \min_{t \in [\delta, 1-\delta]} u(t) = \alpha(u) \geq d_1$$

for all  $t \in [\delta, 1-\delta]$ . Thus, in view of (2.5), we see that

$$\begin{aligned} \alpha(Au) &= \min_{t \in [\delta, 1-\delta]} \left[ \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \right. \\ &\quad \left. + \frac{\alpha\phi_1(t)}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \right] \\ &\geq \min_{t \in [\delta, 1-\delta]} \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \\ &\quad + \frac{\alpha\phi_1(\delta)}{1 - \alpha\phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \\ &\geq \min_{t \in [\delta, 1-\delta]} \int_{\delta}^{1-\delta} G(t, s) p(s) h(s) f(u(s)) ds \\ &\quad + \frac{\alpha\phi_1(\delta)}{1 - \alpha\phi_1(\eta)} \int_{\delta}^{1-\delta} G(\eta, s) p(s) h(s) f(u(s)) ds \\ &> \frac{d_1}{C} \left[ \min_{t \in [\delta, 1-\delta]} \int_{\delta}^{1-\delta} G(t, s) p(s) h(s) ds \right. \\ &\quad \left. + \frac{\alpha\phi_1(\delta)}{1 - \alpha\phi_1(\eta)} \int_{\delta}^{1-\delta} G(\eta, s) p(s) h(s) ds \right] = d_1, \end{aligned}$$

as required.

Finally, we assert that if  $u \in P(\alpha, d_1, c)$  and  $\|Au\| > d_1/\gamma$ , then  $\alpha(Au) > d_1$ .

To see this, suppose  $u \in P(\alpha, d_1, c)$  and  $\|Au\| > d_1/\gamma$ , then, in view of (2.2) and (2.3), we have

$$\begin{aligned}
\alpha(Au) &= \min_{t \in [\delta, 1-\delta]} \left[ \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \right. \\
&\quad \left. + \frac{\alpha \phi_1(t)}{1 - \alpha \phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \right] \\
&\geq \min_{t \in [\delta, 1-\delta]} \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \\
&\quad + \frac{\alpha \phi_1(\delta)}{1 - \alpha \phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \\
&\geq \gamma \left[ \int_0^1 G(s, s) p(s) h(s) f(u(s)) ds \right. \\
&\quad \left. + \frac{\alpha}{1 - \alpha \phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \right] \\
&\geq \gamma \left[ \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \right. \\
&\quad \left. + \frac{\alpha \phi_1(t)}{1 - \alpha \phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \right]
\end{aligned}$$

for  $t \in [0, 1]$ . Thus

$$\begin{aligned}
\alpha(Au) &\geq \gamma \max_{t \in [0, 1]} \left[ \int_0^1 G(t, s) p(s) h(s) f(u(s)) ds \right. \\
&\quad \left. + \frac{\alpha}{1 - \alpha \phi_1(\eta)} \int_0^1 G(\eta, s) p(s) h(s) f(u(s)) ds \phi_1(t) \right] \\
&= \gamma \|Au\| > \gamma \frac{d_1}{\gamma} = d_1.
\end{aligned}$$

To sum up, all the hypotheses of the Leggett–Williams theorem are satisfied by taking  $b_0 = d_1/\gamma$ . Hence  $A$  has at least three fixed points, i.e., the boundary value problem (1.1)–(1.2) has at least three positive solutions  $u$ ,  $v$  and  $w$  such that

$$\|u\| < d_0, \quad d_1 < \min_{t \in [\delta, 1-\delta]} v(t), \quad \|w\| > d_0,$$

and

$$\min_{t \in [\delta, 1-\delta]} w(t) < d_1.$$

The proof is complete.  $\square$

We remark that the condition (H5) in Theorem 2.1 can be replaced by the following condition (H5'):

$$\lim_{u \rightarrow \infty} \frac{f(u)}{u} < \frac{1}{D},$$

which is a special case of (H5).

**Corollary 2.1.** *If the condition (H5) in Theorem 2.1 is replaced by (H5'), then the conclusion of Theorem 2.1 also holds.*

**Proof.** By Theorem 2.1, we only need to prove that (H5') implies that (H5) holds, that is, if (H5') holds, then there exists a number  $c > d_1/\gamma$  such that  $f(u) < c/D$  for  $u \in [0, c]$ .

Suppose on the contrary that for any  $c > d_1/\gamma$ , there exists  $u_c \in [0, c]$  such that  $f(u_c) \geq c/D$ . Hence if we choose  $c_n > d_1/\gamma$  ( $n = 1, 2, \dots$ ) with  $c_n \rightarrow \infty$ , then there exist  $u_n \in [0, c_n]$  such that

$$f(u_n) \geq \frac{c_n}{D}, \quad (2.6)$$

and so

$$\lim_{n \rightarrow \infty} f(u_n) = \infty. \quad (2.7)$$

Since the condition (H5') holds, then there exists  $\tau > 0$  such that

$$f(u) < \frac{u}{D}, \quad u > \tau. \quad (2.8)$$

Hence we have  $u_n \leq \tau$ . Otherwise, if  $u_n > \tau$ , then it follows from (2.8) that

$$f(u_n) < \frac{u_n}{D} \leq \frac{c_n}{D}, \quad (2.9)$$

which contradicts (2.6).

Let

$$M = \max_{u \in [0, \tau]} f(u).$$

Then  $f(u_n) \leq M$  ( $n = 1, 2, \dots$ ), which also contradicts  $\lim_{n \rightarrow \infty} f(u_n) = \infty$ . The proof is complete.  $\square$

**Example 2.1.** Consider the boundary value problem

$$u''(t) - u(t) + f(u(t)) = 0, \quad t \in (0, 1), \quad (2.10)$$

$$u(0) = 0, \quad u(1) = e^{1/2}u(1/2), \quad (2.11)$$

where

$$f(x) = \begin{cases} x^2, & x \in \left[0, \frac{4(e^2-1)(e^{1/2}+1)}{(2e^{5/4}-2e-e^{1/2}+1)(e^{5/4}-e^{3/4})}\right], \\ lx+k, & x \in \left[\frac{4(e^2-1)(e^{1/2}+1)}{(2e^{5/4}-2e-e^{1/2}+1)(e^{5/4}-e^{3/4})}, +\infty\right), \end{cases}$$

$$l = \frac{e+1}{2(e+1-2e^{1/2})(1+e^{3/2}+e^{1/2})},$$

and

$$k = \left( \frac{4(e^2-1)(e^{1/2}+1)}{(2e^{5/4}-2e-e^{1/2}+1)(e^{5/4}-e^{3/4})} \right)^2 - \frac{4(e+1)(e^2-1)(e^{1/2}+1)}{2(e+1-2e^{1/2})(1+e^{3/2}+e^{1/2})(2e^{5/4}-2e-e^{1/2}+1)(e^{5/4}-e^{3/4})}.$$

A simple calculation shows that

$$\phi_1(t) = \frac{e^{1+t} - e^{1-t}}{e^2 - 1}, \quad \phi_2(t) = \frac{e^{2-t} - e^t}{e^2 - 1}, \quad \phi_1'(0) = \frac{2e}{e^2 - 1}, \quad \text{and} \\ p(t) \equiv 1.$$

Take  $\delta = 1/4$ ; then it is not difficult to obtain that

$$D = \frac{(e+1-2e^{1/2})(1+e^{3/2}+e^{1/2})}{e+1}, \quad C = \frac{2e^{5/4}-2e-e^{1/2}+1}{2(e^{1/2}+1)}, \quad \text{and} \\ \gamma = \frac{e^{5/4}-e^{3/4}}{e^2-1}.$$

Thus, if we choose

$$d_0 = \frac{e+1}{2(e+1-2e^{1/2})(1+e^{3/2}+e^{1/2})} \quad \text{and} \quad d_1 = \frac{4(e^{1/2}+1)}{2e^{5/4}-2e-e^{1/2}+1},$$

then the conditions (2.4), (2.5) and (H5') are fulfilled. It follows from Corollary 2.1 that the boundary value problem (2.10)–(2.11) has at least three positive solutions.

## References

- [1] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, New York, 1985.
- [2] L. Erbe, H. Wang, On the existence of positive solutions of ordinary differential equations, *Proc. Amer. Math. Soc.* 120 (1994) 743–748.
- [3] D. Guo, V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, Academic Press, San Diego, 1988.
- [4] C.P. Gupta, Solvability of a three-point nonlinear boundary value problems for a second order ordinary differential equation, *J. Math. Anal. Appl.* 168 (1992) 540–551.
- [5] C.P. Gupta, A sharp condition for the solvability of a three-point second order boundary value problem, *J. Math. Anal. Appl.* 205 (1997) 579–586.
- [6] C.P. Gupta, S. Trofimchuk, Existence of a solution to a three-point boundary value problem and the spectral radius of a related linear operator, *Nonlinear Anal.* 34 (1998) 498–507.
- [7] R.W. Leggett, L.R. Williams, Multiple positive fixed points of nonlinear operators on ordered Banach spaces, *Indiana Univ. Math. J.* 28 (1979) 673–688.
- [8] R. Ma, Existence theorems for a second order three-point boundary value problem, *J. Math. Anal. Appl.* 212 (1997) 430–442.



- [9] R. Ma, Positive solutions of a nonlinear three-point boundary value problem, *Electron. J. Differential Equations* 34 (1999) 1–8.
- [10] R. Ma, H. Wang, Positive solutions of nonlinear three-point boundary value problems, *J. Math. Anal. Appl.* 279 (2003) 216–227.
- [11] J.P. Sun, H.F. Huo, W.T. Li, Positive solutions to singular nonlinear three point boundary value problems, *Dynamic Systems and Applications*, in press.
- [12] H. Wang, On the existence of positive solution for semilinear elliptic equations in the annulus, *J. Differential Equations* 109 (1994) 1–7.